

ASYMPTOTICALLY EFFICIENT ESTIMATION OF LINEAR FUNCTIONALS IN INVERSE REGRESSION MODELS

CHRIS A.J. KLAASSEN, EUN-JOO LEE, AND FRITS H. RUYMGAART*

ABSTRACT. In this paper we will discuss a procedure to improve the usual estimator of a linear functional of the unknown regression function in inverse nonparametric regression models. In Klaassen *et al.* (2001) it has been proved that this traditional estimator is not asymptotically efficient (in the sense of the Hájek - Le Cam convolution theorem) except, possibly, when the error distribution is normal. Since this estimator, however, is still root-n consistent a procedure in Bickel *et al.* (1993) applies to construct a modification which is asymptotically efficient. A self-contained proof of the asymptotic efficiency is included.

1. Introduction

In the nonparametric regression model with either direct or indirect observations, both the parameter of actual interest (the unknown regression function) and the nuisance parameter (the error density that may also be unknown) are infinite dimensional. In this paper we consider the problem of asymptotically efficient (AE) estimation of a linear functional of the regression function. In other words, our aim is to construct estimators that are asymptotically normal with smallest possible variance. Linear functionals are of independent interest and often studied in the literature (Ibragimov & Hasminskii (1984), Goldenshluger & Pereverzev (2000)). They are also important because the Fourier coefficients of an expansion of the regression function in an orthonormal basis are linear functionals. Below we will briefly return to the latter aspect.

Suitable estimators of the regression function naturally yield estimators of the linear functional by substitution in the inner product representing this functional. In van Rooij *et al.* (1999) it has been proved that, when a linear functional of an indirectly sampled density is to be estimated, substitution of a suitable density estimator produces an AE estimator of the functional. Even when the error density is known but arbitrary, the situation to be considered here, AE estimation of a linear functional of the regression function turns out to be an essentially harder problem. Klaassen *et al.* (2001) have shown that substitution of the usual type of regression function estimator does not produce an AE estimator of the functional, except possibly when the error distribution is normal. Another, somewhat simpler natural estimator could be easily proposed for linear functionals, but this estimator is equivalent with and in certain cases even identical to the plug-in estimator and hence not AE either. It follows in particular that the usual orthonormal series

1991 *Mathematics Subject Classification.* Primary 62G08, 62G20; Secondary 45R05.

Key words and phrases. Noisy integral equation, indirect nonparametric regression, linear functional, asymptotic efficiency, improving a root-n consistent estimator.

*Research partly supported by National Science Foundation grant DMS-0203942.

type estimator of the regression function is obtained from estimators of the Fourier coefficients that are not AE in the above sense. Nevertheless such estimators do in general attain the best convergence rate for the mean integrated squared error.

In this paper we will focus on the question of how to improve the plug-in estimator of a linear functional. The plug-in-method is a simple device to construct an estimator of just about any functional. We will see, moreover, that an estimator of the regression function will also be needed in the improvement procedure. Although not AE, the plug-in estimator is $\sqrt{n}$ -consistent. This means that the method to construct an AE estimator given a $\sqrt{n}$ -consistent one, described in great generality in Bickel *et al.* (1993) for infinite dimensional parameters, applies. See also Pfanzagl (1994) for improving $\sqrt{n}$ -consistent estimators in models with finite dimensional parameter. In principle our result would fit into the general theory as described in particular in Chapter 7 of Bickel *et al.* (1993). We prefer, however, to provide a self-contained and independent derivation in this paper. On the one hand the indirect regression model is sufficiently specific to allow explicit calculations, on the other hand it is of sufficient importance to warrant such an effort. The question whether employing improved estimators of the Fourier coefficients in a series type estimator of the regression function itself would improve this estimator in a certain way might be of some interest. Although this cannot be true for the rate, we suspect that improvement will hold true at the level of constants. Investigating this matter is beyond the scope of this paper. In Section 6, however, we will briefly comment on this.

In order to construct an AE estimator one needs to compute the efficient influence function. According to van der Vaart (1998) this is the projection of the gradient of the functional onto the tangent space to the model. See Section 3 for the details. Statement and proof that the improved estimator is AE can be found in Section 4. Indirect nonparametric regression occurs in a wide variety of practical situations, like Wicksell's unfolding problem in stereology, geological prospecting, computer tomography in imaging, just to mention a few (O'Sullivan (1986), Kress (1989), Kirsch (1996)). Let us introduce another example (see also Section 6.1).

Example. Can you see the weight of a cable? To answer this question, a paraphrase of the title of Kac's famous 1966 paper, let us first observe that the shape of a cable suspended at its endpoints with coordinates (0,0) and (1,0) is given by the differential equation

$$(1.1) \quad -\frac{d^2 g(t)}{dt^2} = f(t), \quad 0 \leq t \leq 1, \quad g(0) = g(1) = 0.$$

Apart from the sign the source term f represents the load per horizontal distance and g the shape. The problem is to estimate a linear functional $\int_0^1 f(t)\varphi(t) dt$, for suitable φ , like for instance the total weight $\int_0^1 f(t) dt$ of the cable ($\varphi \equiv 1$); for this special case, however, see the remark in Section 6.1. Estimation may be performed by first recovering the weight distribution or source term from the data $(X_1, Y_1), \dots, (X_n, Y_n)$ that are independent copies of (X, Y) , where

$$(1.2) \quad Y = g(X) + \varepsilon.$$

We will assume that the design variable X has the Uniform (0,1) distribution and is independent of the error variable ε . The latter has also zero mean, finite variance, and arbitrary density ψ that is supposed to be known.

Using the Green's function the differential equation can be rewritten in the form

$$(1.3) \quad g(x) = \int_0^1 K(x, t) f(t) dt = (Kf)(x), \quad 0 \leq x \leq 1,$$

where $K : L^2([0, 1]) \rightarrow L^2([0, 1])$ turns out to be compact and strictly positive Hermitian with Brownian bridge covariance kernel

$$(1.4) \quad K(x, t) = x \wedge t - xt, \quad 0 \leq x \leq 1, \quad 0 \leq t \leq 1.$$

The operator has strictly positive eigenvalues

$$(1.5) \quad \rho_k = \left\{ \frac{1}{\pi(k+1)} \right\}^2, \quad k = 0, 1, \dots,$$

with corresponding orthonormal basis of eigenfunctions

$$(1.6) \quad \varphi_k(x) = \sqrt{2} \sin(k+1)\pi x, \quad 0 \leq x \leq 1, \quad k = 0, 1, \dots$$

The general theory below is tailored to but slightly more general than the compact case. This generalization ensures that the noncompact identity operator, which yields the usual direct regression, is included as a special case.

2. The model, the problem, and assumptions

Let $(\mathbf{X}, \mathcal{X}, \mu)$ and $(\mathbf{Z}, \mathcal{Z}, \nu)$ be measurable spaces with $L^2(\mu)$ and $L^2(\nu)$ real separable Hilbert spaces. We are given a bounded, injective linear operator $K : L^2(\nu) \rightarrow L^2(\mu)$. In the random design case, to be considered here, we observe a random sample $(X_1, Y_1), \dots, (X_n, Y_n)$ of independent copies of a random element (X, Y) , where

$$(2.1) \quad Y = (Kf)(X) + \varepsilon = g(X) + \varepsilon, \quad f \in L^2(\nu).$$

The indirectly observed regression or input function f on $(\mathbf{Z}, \mathcal{Z})$ is unknown. We will assume that

$$(2.2) \quad \mu(\mathbf{X}) = 1, \text{ and } X =_d \text{Uniform}(\mathbf{X}).$$

The error variable ε is independent of the design X with known but arbitrary density ψ with respect to Lebesgue measure. We will assume that

$$(2.3) \quad \mathbf{E}\varepsilon = 0, \quad \mathbf{E}\varepsilon^2 = \sigma^2.$$

Under these assumptions it is readily verified that (X, Y) has density

$$(2.4) \quad p_f(x, y) = \psi(y - (Kf)(x)), \quad (x, y) \in \mathbf{X} \times \mathbb{R}, \quad f \in L^2(\nu).$$

A star attached to an operator denotes its adjoint. The operator $R = (K^*K)^{\frac{1}{2}} : L^2(\nu) \rightarrow L^2(\nu)$, then, is strictly positive Hermitian. We will assume that there exists an orthonormal basis for $L^2(\nu)$ consisting of eigenfunctions $\varphi_0, \varphi_1, \dots$ of the operator R with corresponding eigenvalues $\rho_0, \rho_1, \dots > 0$ satisfying

$$(2.5) \quad \sup_{k \geq 0} \rho_k < \infty.$$

On the one hand compact operators R are included, since for those a basis exists with eigenvalues satisfying $\rho_k \downarrow 0$, as $k \rightarrow \infty$ (Debnath & Mikusiński (1999)). On the other hand the direct model with $K = I$ and hence $R = I$, where I is the identity operator, is included as well. The operator I is not compact but

satisfies the condition above for any basis, with $\rho_k = 1$ for all k . According to the polar decomposition (Riesz & Nagy (1990)) there exists a partial isometry $V : L^2(\nu) \rightarrow L^2(\mu)$ such that $K = VR$ and $K^* = RV^*$. It should be noted that V^*V is the identity on the range of R . Let us write

$$(2.6) \quad \varphi_{V,k} = V\varphi_k,$$

and observe that the $\varphi_{V,k}$ are orthonormal in $L^2(\mu)$.

The problem to be considered here is estimation of a linear functional $f \mapsto \langle f, \varphi \rangle$, $f \in L^2(\nu)$, for some given $\varphi \in L^2(\nu)$ with $\|\varphi\| = 1$. In view of $\langle f, \varphi \rangle = \sum_{k \geq 0} \langle f, \varphi_k \rangle \langle \varphi, \varphi_k \rangle$ it seems plausible that estimation of the special linear functionals defined by the Fourier coefficients

$$(2.7) \quad f_k = \langle f, \varphi_k \rangle$$

might suffice. This is in fact true under an extra condition, and some details can be found in Section 5. Hence we will focus on estimating an arbitrary Fourier coefficient. As a generic example let us consider the functional

$$(2.8) \quad f \mapsto f_0 = \langle f, \varphi_0 \rangle, f \in L^2(\nu),$$

where φ_0 is the first basis element. It is useful to observe

$$(2.9) \quad f_k = \langle f, \varphi_k \rangle = \langle f, K^*VR^{-1}\varphi_k \rangle = \langle Kf, VR^{-1}\varphi_k \rangle = \frac{1}{\rho_k} \langle Kf, \varphi_{V,k} \rangle.$$

The assumptions below will be briefly discussed in Section 6. The basis elements are supposed to satisfy the uniform boundedness conditions

$$(2.10) \quad \sup_{z \in \mathbf{Z}, k \geq 0} |\varphi_k(z)| < \infty$$

and

$$(2.11) \quad \sup_{x \in \mathbf{X}, k \geq 0} |\varphi_{V,k}(x)| < \infty.$$

Regarding the input function f it will be assumed that there exists a sequence $(m(n))_{n \geq 1}$ satisfying

$$(2.12) \quad m = m(n) \rightarrow \infty \text{ and } \frac{m}{\sqrt{n}} \rightarrow 0, \text{ as } n \rightarrow \infty,$$

for which

$$(2.13) \quad \sqrt{n} \sum_{k > m} f_k^2 \rightarrow 0, \text{ as } n \rightarrow \infty,$$

$$(2.14) \quad \sum_{k > m} |f_k| \rightarrow 0, \text{ as } n \rightarrow \infty.$$

The error density ψ is supposed to be twice differentiable, i.e.,

$$(2.15) \quad \psi'' \text{ exists on } \mathbb{R}.$$

Denoting the score function for location by

$$(2.16) \quad \Lambda = -\frac{\psi'}{\psi},$$

we will also need that

$$(2.17) \quad \Lambda' \text{ and } \Lambda'' \text{ exist and are bounded on } \mathbb{R},$$

with finite Fisher information

$$(2.18) \quad \mathbf{E}\Lambda^2(\varepsilon) = \mathbf{E}\Lambda'(\varepsilon) = \int_{-\infty}^{\infty} \left(\frac{\psi'}{\psi}\right)^2 (y)\psi(y) dy = I(\psi) < \infty.$$

For the first equality in (2.18) we need (2.15).

3. Construction of an asymptotically efficient estimator

The input function $f \in L^2(\nu)$ has the L^2 - expansion

$$(3.1) \quad f(z) = \sum_{k \geq 0} f_k \varphi_k(z), \quad z \in \mathbf{Z},$$

in the orthonormal basis of eigenfunctions, and the usual estimator of the regression function in this context is given by

$$(3.2) \quad \hat{f}(z) = \hat{f}_{(m)}(z) = \sum_{k \leq m} \hat{f}_k \varphi_k(z), \quad z \in \mathbf{Z},$$

for suitable $m = m(n) \rightarrow \infty$, as $n \rightarrow \infty$, where $\hat{f}_k$ is the estimator of f_k (see (2.1) and (2.9)) given by

$$(3.3) \quad \hat{f}_k = \frac{1}{n} \sum_{i=1}^n \frac{1}{\rho_k} Y_i \varphi_{V,k}(X_i).$$

See, for instance, Johnstone & Silverman (1990). Since we are estimating f_0 in (2.8), a plug-in estimator simply equals

$$(3.4) \quad \langle \hat{f}, \varphi_0 \rangle = \hat{f}_0,$$

as given by (3.3) for $k = 0$.

The estimators $\hat{f}_k (k = 0, 1, \dots)$ have some desirable properties. Because of

$$(3.5) \quad \begin{aligned} \mathbf{E}\hat{f}_k &= \frac{1}{\rho_k} \mathbf{E}Y \varphi_{V,k}(X) = \frac{1}{\rho_k} \mathbf{E}(Kf)(X) \varphi_{V,k}(X) \\ &= \frac{1}{\rho_k} \langle Kf, V \varphi_k \rangle = \frac{1}{\rho_k} \langle f, RV^* V \varphi_k \rangle \\ &= \frac{1}{\rho_k} \rho_k \langle f, \varphi_k \rangle = f_k \end{aligned}$$

they are unbiased, but $\hat{f} = \hat{f}_{(m)}$ is not; let us write

$$(3.6) \quad f_{(m)}(z) = \mathbf{E}\hat{f}_{(m)}(z) = \sum_{k \leq m} f_k \varphi_k(z), \quad z \in \mathbf{Z}.$$

Furthermore we have

$$(3.7) \quad \begin{aligned} \mathbf{E}(\hat{f}_k - f_k)^2 &= \mathbf{Var} \hat{f}_k = \frac{1}{n} \mathbf{Var} \frac{1}{\rho_k} Y \varphi_{V,k}(X) \\ &\leq \frac{1}{n} \left(\frac{1}{\rho_k}\right)^2 \mathbf{E}Y^2 \varphi_{V,k}^2(X) \leq C \frac{1}{n} \left(\frac{1}{\rho_k}\right)^2, \end{aligned}$$

where $0 < C < \infty$ will throughout be used as a generic constant that will not depend on n or k and that here can be taken equal to

$$(3.8) \quad C = (\sup_{x \in \mathbf{X}, k \geq 0} |\varphi_{V,k}^2(x)|) \cdot (\|Kf\|^2 + \sigma^2),$$

by assumption (2.11).

The central limit theorem yields at once the asymptotic normality of the empirical Fourier coefficients. In particular we have

$$(3.9) \quad \sqrt{n}(\hat{f}_0 - f_0) \rightarrow_d \text{Normal}(0, \sigma_0^2(f)), \text{ as } n \rightarrow \infty,$$

where (cf. (2.1), (2.3), and (2.6))

$$(3.10) \quad \begin{aligned} \sigma_0^2(f) &= \left(\frac{1}{\rho_0}\right)^2 \mathbf{Var}(Y\varphi_{V,0}(X)) \\ &\geq \left(\frac{1}{\rho_0}\right)^2 \mathbf{E} \mathbf{Var}(Y\varphi_{V,0}(X)|X) \\ &= \left(\frac{\sigma}{\rho_0}\right)^2 \mathbf{E} \varphi_{V,0}^2(X) = \left(\frac{\sigma}{\rho_0}\right)^2 \end{aligned}$$

holds with a strict inequality, unless $\mathbf{Var}(\mathbf{E}(Y\varphi_{V,0}(X)|X))$ vanishes, i.e., unless $(Kf)(X)\varphi_{V,0}(X)$ is degenerate.

It has already been observed in Klaassen *et al.* (2001) that $\sigma_0^2(f)$ is in general strictly larger than the optimal variance according to the Hájek - Le Cam convolution theorem (van der Vaart (1998)). In other words the estimator $\hat{f}_0$ is not asymptotically efficient, but it is $\sqrt{n}$ -consistent. Such estimators can be improved. In order to do so we first need to briefly review some results from Klaassen *et al.* (2001).

Since we assume ψ to be known, in terms of square roots of the densities the model is $\mathcal{S} = \{s_f, f \in L^2(\nu)\}$, with $s_f = \sqrt{p_f}$ and p_f as in (2.4). The tangent space at $f \in L^2(\nu)$ to this model is given by

$$(3.11) \quad \dot{\mathcal{S}}_f = [\dot{s}_{f,h}, h \in \mathfrak{R}_K] = \{\dot{s}_{f,h}, h \in \overline{\mathfrak{R}_K}\} \subset L^2(\mu \times \lambda),$$

where λ is Lebesgue measure on $\mathbb{R}$, $\mathfrak{R}_K$ is the range of K , $\overline{\mathfrak{R}_K}$ its closure, and

$$(3.12) \quad \dot{s}_{f,h}(x, y) = -\frac{\psi'(y - (Kf)(x))}{2s_f(x, y)} h(x), \quad x \in \mathbf{X}, y \in \mathbb{R}.$$

The gradient of the functional $f \mapsto T(s_f) = \langle f, \varphi_0 \rangle$ at f is given by (cf. (2.9))

$$(3.13) \quad \dot{T}_f(x, y) = \frac{2}{\rho_0}(y - (Kf)(x))\varphi_{V,0}(x)s_f(x, y), \quad x \in \mathbf{X}, y \in \mathbb{R}.$$

Let $\tilde{T}_f$ denote the projection in $L^2(\mu \times \lambda)$ of $\dot{T}_f$ onto $\dot{\mathcal{S}}_f$. Then the optimal variance mentioned in the preceding paragraph equals

$$(3.14) \quad \tilde{\sigma}_0^2(f) = \frac{1}{4}\|\tilde{T}_f\|^2.$$

For these results see Klaassen *et al.* (2001).

In order to construct an estimator with limiting normal distribution having the variance in (3.14) we first need to explicitly compute $\tilde{T}_f$. Because $\tilde{T}_f$ has to belong to $\dot{\mathcal{S}}_f$ it is of the form

$$(3.15) \quad \tilde{T}_f(x, y) = -\frac{\psi'(y - (Kf)(x))}{2s_f(x, y)} \tilde{h}(x), \quad x \in \mathbf{X}, y \in \mathbb{R},$$

for some $\tilde{h} \in \overline{\mathfrak{R}_K}$, and because of $\dot{T}_f - \tilde{T}_f \perp \dot{\mathcal{S}}_f$ we must have (see also (3.12))

$$(3.16) \quad \langle \dot{T}_f - \tilde{T}_f, \dot{\mathcal{S}}_{f,h} \rangle = \iint \left\{ \frac{2}{\rho_0} (y - (Kf)(x)) \varphi_{V,0}(x) s_f(x, y) + \frac{\psi'(y - (Kf)(x))}{2s_f(x, y)} \tilde{h}(x) \right\} \\ \times \left\{ \frac{\psi'(y - (Kf)(x))}{2s_f(x, y)} h(x) \right\} d\mu(x) dy = 0,$$

for all $h \in \overline{\mathfrak{R}_K}$. It follows from (2.3) and (2.15) that $\int_{-\infty}^{\infty} y\psi'(y) dy = -1$ holds; cf. Lemma I.2.4.b of Hájek and Šidák (1967). Exploiting this fact and using the notation $I(\psi)$ defined in (2.18) straightforward integration shows that (3.16) entails

$$(3.17) \quad \langle I(\psi)\tilde{h} - \frac{4}{\rho_0} \varphi_{V,0}, h \rangle = 0, \text{ for all } h \in \overline{\mathfrak{R}_K}.$$

It is obvious that $\frac{1}{\rho_0} \varphi_{V,0} \in \mathfrak{R}_K$. Since $\tilde{h} \in \overline{\mathfrak{R}_K}$ it follows from (3.17) that

$$(3.18) \quad \tilde{h}(x) = \frac{4}{I(\psi)\rho_0} \varphi_{V,0}(x), \quad x \in \mathbf{X}.$$

Combination with (3.14) yields

$$(3.19) \quad \tilde{\sigma}_0^2(f) = \frac{1}{4} \|\tilde{T}_f\|^2 = \frac{1}{\rho_0^2 I(\psi)},$$

for the value of the optimal variance. Note that

$$1 = \left(\int_{-\infty}^{\infty} y\psi'(y) dy \right)^2 \leq \sigma^2 I(\psi)$$

holds by the Cramér - Rao inequality with equality if and only if $\psi'(Y)/\psi(Y)$ is linear in Y a.s. under ψ , i.e. if and only if ψ is a normal density.

Consequently, it is immediate from (3.10) that the string of inequalities

$$(3.20) \quad \sigma_0^2(f) \geq \left(\frac{1}{\rho_0} \right)^2 \sigma^2 \geq \left(\frac{1}{\rho_0} \right)^2 \frac{1}{I(\psi)} = \tilde{\sigma}_0^2(f)$$

holds. For nonnormal densities ψ the second inequality is strict, and the estimator $\hat{f}_0$ then turns out not to be asymptotically efficient. If ψ is normal, we have $I(\psi) = 1/\sigma^2$, so that the second inequality in (3.20) is an equality. However, asymptotic efficiency of $\hat{f}_0$ remains impossible, since the first inequality in (3.20) cannot be an equality for all f , as argued in (3.10). See Klaassen *et al.* (2001) for some details.

We are now in a position to construct the improved estimator. According to van der Vaart (1998) define the efficient influence function by

$$(3.21) \quad \tilde{T}_f(x, y) \cdot \frac{1}{2s_f(x, y)} = -\frac{1}{I(\psi)} \frac{\psi'(y - (Kf)(x))}{\psi(y - (Kf)(x))} \frac{1}{\rho_0} \varphi_{V,0}(x) \\ = \frac{1}{I(\psi)} \Lambda(y - (Kf)(x)) \frac{1}{\rho_0} \varphi_{V,0}(x), \quad x \in \mathbf{X}, y \in \mathbb{R},$$

using the notation Λ introduced in (2.16). Following a procedure in Bickel *et al.* (1993, Chapter 7) let us now introduce the estimator

$$(3.22) \quad \hat{f}_0 = \hat{f}_0 + \frac{1}{n} \sum_{i=1}^n \frac{1}{\rho_0 I(\psi)} \Lambda(Y_i - (K\hat{f}_{(m)})(X_i)) \varphi_{V,0}(X_i),$$

where $\hat{f}_{(m)} = \hat{f}$ is defined in (3.2). We will see in the next section that this turns out to be an asymptotically efficient estimator of f_0 .

4. The main theorem

Theorem. *Suppose that all the assumptions listed in Section 2 are fulfilled. Then $\hat{f}_0$ defined in (3.22) is an asymptotically efficient estimator of f_0 , i.e.*

$$(4.1) \quad \sqrt{n}(\hat{f}_0 - f_0) \rightarrow_d \text{Normal} \left(0, \frac{1}{\rho_0^2 I(\psi)} \right), \text{ as } n \rightarrow \infty,$$

where the variance in the normal distribution is optimal.

Proof. A Taylor expansion yields

$$(4.2) \quad \sqrt{n}(\hat{f}_0 - f_0) = A_n + Q_n + R_n$$

with

$$(4.3) \quad A_n = \frac{1}{\sqrt{n}} \sum_{i=1}^n \frac{1}{\rho_0 I(\psi)} \Lambda(\varepsilon_i) \varphi_{V,0}(X_i),$$

$$(4.4) \quad Q_n = \sqrt{n}[\langle \hat{f} - f, \varphi_0 \rangle - \frac{1}{n} \sum_{i=1}^n \frac{1}{\rho_0 I(\psi)} \{(K\hat{f})(X_i) - (Kf)(X_i)\} \Lambda'(\varepsilon_i) \varphi_{V,0}(X_i)],$$

$$(4.5) \quad R_n = \frac{1}{2\sqrt{n}} \sum_{i=1}^n \frac{1}{\rho_0 I(\psi)} \{(K\hat{f})(X_i) - (Kf)(X_i)\}^2 \Lambda''(\tilde{\varepsilon}_i) \varphi_{V,0}(X_i).$$

In (4.5), $\tilde{\varepsilon}_i$ is a random variable between $Y_i - (Kf)(X_i) = \varepsilon_i$ and $Y_i - (K\hat{f})(X_i)$.

It follows from (2.11) that

$$(4.6) \quad \sup_{x \in \mathbf{X}} |\varphi_{V,0}(x)| < \infty.$$

Together with (2.17) we find (C generic !)

$$(4.7) \quad \begin{aligned} |R_n| &\leq \frac{C}{\sqrt{n}} \sum_{i=1}^n \{(K\hat{f})(X_i) - (Kf)(X_i)\}^2 \\ &= \frac{C}{\sqrt{n}} \sum_{i=1}^n \left\{ \sum_{k \leq m} (\hat{f}_k - f_k) \rho_k \varphi_{V,k}(X_i) - \sum_{k > m} f_k \rho_k \varphi_{V,k}(X_i) \right\}^2 \\ &\leq \frac{C}{\sqrt{n}} \sum_{k \leq m} \sum_{l \leq m} (\hat{f}_k - f_k)(\hat{f}_l - f_l) \rho_k \rho_l \left\{ \sum_{i=1}^n \varphi_{V,k}(X_i) \varphi_{V,l}(X_i) \right\} \\ &\quad + \frac{C}{\sqrt{n}} \sum_{k > m} \sum_{l > m} f_k f_l \rho_k \rho_l \left\{ \sum_{i=1}^n \varphi_{V,k}(X_i) \varphi_{V,l}(X_i) \right\} \\ &= R_{n1} + R_{n2}. \end{aligned}$$

It clearly suffices to show that $\mathbf{E}R_{nj} \rightarrow 0$, as $n \rightarrow \infty$, for $j = 1, 2$, in order to obtain $R_n \rightarrow_p 0$, as $n \rightarrow \infty$.

Regarding the first term we see that

$$(4.8) \quad \mathbf{E}R_{n1} = \frac{C}{n^{5/2}} \sum_{k \leq m} \sum_{l \leq m} \left[\sum_{\alpha=1}^n \sum_{\beta=1}^n \sum_{i=1}^n \mathbf{E}\{Y_\alpha \varphi_{V,k}(X_\alpha) - f_k \rho_k\} \right. \\ \left. \times \{Y_\beta \varphi_{V,l}(X_\beta) - f_l \rho_l\} \{\varphi_{V,k}(X_i) \varphi_{V,l}(X_i)\} \right].$$

By decomposing $\sum \sum \sum_{\alpha, \beta, i} = \sum \sum \sum_{\alpha \neq \beta \neq i} + \sum \sum_{\alpha \neq \beta = i} + \sum \sum_{\beta \neq \alpha = i} + \sum \sum_{\alpha = \beta \neq i} + \sum_{\alpha = \beta = i}$, and by realizing that the random variables labeled with α and β are centered at 0 and that the $\varphi_{V,k}$ are orthonormal, we arrive at (δ_{kl} is Kronecker's delta)

$$(4.9) \quad 0 \leq \mathbf{E}R_{n1} \\ = \frac{C}{n^2 \sqrt{n}} \sum_{k \leq m} \sum_{l \leq m} \{n(n-1) \delta_{kl} \mathbf{Cov}(Y \varphi_{V,k}(X), Y \varphi_{V,l}(X)) \\ + n \mathbf{E}(Y \varphi_{V,k}(X) - f_k \rho_k)(Y \varphi_{V,l}(X) - f_l \rho_l) \varphi_{V,k}(X) \varphi_{V,l}(X)\} \\ \leq \frac{C}{n^2 \sqrt{n}} [n(n-1) \sum_{k \leq m} \mathbf{E}Y^2 \varphi_{V,k}^2(X) \\ + n \sum_{k \leq m} \sum_{l \leq m} \{\mathbf{E}Y^2 \varphi_{V,k}^2(X) \mathbf{E}Y^2 \varphi_{V,l}^2(X)\}^{1/2}] \\ \leq C \left(\frac{m}{\sqrt{n}} + \frac{m^2}{n\sqrt{n}} \right) \rightarrow 0, \text{ as } n \rightarrow \infty.$$

Here we have used (2.11) and (2.12). See also the calculation in (3.7).

Next let us observe that

$$(4.10) \quad 0 \leq \mathbf{E}R_{n2} = \frac{C}{\sqrt{n}} \sum_{k > m} \sum_{l > m} n f_k f_l \rho_k \rho_l \delta_{kl} \\ \leq C \sqrt{n} \sum_{k > m} f_k^2 \rho_k^2 \\ \leq C \sqrt{n} \sum_{k > m} f_k^2 \rightarrow 0, \text{ as } n \rightarrow \infty,$$

by assumption (2.13). This settles the asymptotic negligibility of R_n .

For brevity let us introduce

$$(4.11) \quad \bar{U}_k = \frac{1}{n} \sum_{i=1}^n \left\{ \langle \varphi_0, \varphi_k \rangle - \frac{1}{\rho_0 I(\psi)} \rho_k \varphi_{V,k}(X_i) \Lambda'(\varepsilon_i) \varphi_{V,0}(X_i) \right\},$$

and note that $\bar{U}_k$ is an average of i.i.d. random variables with zero mean. Indeed we have (cf.(2.18))

$$(4.12) \quad \langle \varphi_0, \varphi_k \rangle - \frac{1}{\rho_0 I(\psi)} \rho_k \mathbf{E} \varphi_{V,k}(X) \Lambda'(\varepsilon) \varphi_{V,0}(X) \\ = \delta_{0k} - \frac{1}{\rho_0 I(\psi)} \rho_k I(\psi) \delta_{k0} = 0, \text{ for all } k.$$

Because of

$$(4.13) \quad \begin{aligned} Q_n &= \sqrt{n} \sum_{k \leq m} (\hat{f}_k - f_k) \bar{U}_k - \sqrt{n} \sum_{k > m} f_k \bar{U}_k \\ &= Q_{n1} + Q_{n2}, \end{aligned}$$

in order to show that $Q_n \rightarrow_p 0$, as $n \rightarrow \infty$, it suffices to prove $\mathbf{E}|Q_{n1}| \rightarrow 0$ and $\mathbf{E}|Q_{n2}| \rightarrow 0$.

In order to deal with Q_{n1} let us first note that

$$(4.14) \quad \begin{aligned} \mathbf{E}(\hat{f}_k - f_k)^2 \bar{U}_k^2 &= \\ &= \frac{1}{n^4} \sum_{i=1}^n \sum_{j=1}^n \sum_{\alpha=1}^n \sum_{\beta=1}^n \mathbf{E}\{Y_i \varphi_{V,k}(X_i) - \rho_k f_k\} \{Y_j \varphi_{V,k}(X_j) - \rho_k f_k\} \\ &\quad \times \left\{ \langle \varphi_0, \varphi_k \rangle / \rho_k - \frac{1}{\rho_0 I(\psi)} \varphi_{V,k}(X_\alpha) \Lambda'(\varepsilon_\alpha) \varphi_{V,0}(X_\alpha) \right\} \\ &\quad \times \left\{ \langle \varphi_0, \varphi_k \rangle / \rho_k - \frac{1}{\rho_0 I(\psi)} \varphi_{V,k}(X_\beta) \Lambda'(\varepsilon_\beta) \varphi_{V,0}(X_\beta) \right\}. \end{aligned}$$

Since each factor has zero expectation the only terms that contribute are those with all 4 indices equal and those with 2 pairs of indices (but not all 4) equal. By (2.11), (2.17), and (3.7) the contribution of the first group is seen to be bounded by $C \cdot n$ and the second group is seen to yield a contribution bounded by $C \cdot n^2$. This entails

$$(4.15) \quad \mathbf{E}(\hat{f}_k - f_k)^2 \bar{U}_k^2 \leq \frac{1}{n^4} (C \cdot n + C \cdot n^2) \leq \frac{C}{n^2},$$

and hence, by applying Schwarz's inequality,

$$(4.16) \quad \begin{aligned} \mathbf{E}|Q_{n1}| &\leq \sqrt{n} \sum_{k \leq m} \{\mathbf{E}(\hat{f}_k - f_k)^2 \bar{U}_k^2\}^{1/2} \\ &\leq \sqrt{nm} \frac{C}{n} = C \frac{m}{\sqrt{n}} \rightarrow 0, \text{ as } n \rightarrow \infty, \end{aligned}$$

by assumption (2.12).

For Q_{n2} observe that,

$$(4.17) \quad \begin{aligned} \mathbf{E} \bar{U}_k^2 &= \frac{1}{n^2} \sum_{i=1}^n \sum_{j=1}^n \mathbf{E} \left\{ \langle \varphi_0, \varphi_k \rangle - \frac{1}{\rho_0 I(\psi)} \rho_k \varphi_{V,k}(X_i) \Lambda'(\varepsilon_i) \varphi_{V,0}(X_i) \right\} \\ &\quad \times \left\{ \langle \varphi_0, \varphi_k \rangle - \frac{1}{\rho_0 I(\psi)} \rho_k \varphi_{V,k}(X_j) \Lambda'(\varepsilon_j) \varphi_{V,0}(X_j) \right\} \\ &\leq \frac{C}{n} \left(\frac{\rho_k}{\rho_0} \right)^2 \mathbf{E} \varphi_{V,k}^2(X) (\Lambda')^2(\varepsilon) \varphi_{V,0}^2(X) \\ &\leq \frac{C}{n} \rho_k^2. \end{aligned}$$

This yields, again by the Schwarz inequality

$$(4.18) \quad \begin{aligned} \mathbf{E}|Q_{n2}| &\leq \sqrt{n} \sum_{k > m} |f_k| \{\mathbf{E} \bar{U}_k^2\}^{1/2} \\ &\leq C \sum_{k > m} |f_k| \rho_k \rightarrow 0, \text{ as } n \rightarrow \infty, \end{aligned}$$

by assumptions (2.5) and (2.14).

Finally let us consider A_n in (4.3). Since the terms are i.i.d. with

$$(4.19) \quad \mathbf{E} \frac{1}{\rho_0 I(\psi)} \Lambda(\varepsilon) \varphi_{V,0}(X) = 0,$$

$$(4.20) \quad \begin{aligned} \mathbf{Var} \frac{1}{\rho_0 I(\psi)} \Lambda(\varepsilon) \varphi_{V,0}(X) &= \left(\frac{1}{\rho_0 I(\psi)} \right)^2 \mathbf{E} \Lambda^2(\varepsilon) \mathbf{E} \varphi_{V,0}^2(X) \\ &= \frac{1}{\rho_0^2 I(\psi)}, \end{aligned}$$

the central limit theorem entails at once that

$$(4.21) \quad A_n \rightarrow_d \text{Normal} \left(0, \frac{1}{\rho_0^2 I(\psi)} \right), \text{ as } n \rightarrow \infty.$$

Because we have seen that $R_n \rightarrow_p 0$ and $Q_n \rightarrow_p 0$, this is also the limiting distribution of the expression on the left in (4.2), as was to be shown.

5. Estimating an arbitrary linear functional

In this section we want to consider the problem of estimating the linear functional $f_\varphi = \langle f, \varphi \rangle$, for an arbitrary $\varphi \in L^2(\nu)$ with $\|\varphi\| = 1$. Since $f_\varphi = \sum_{k \geq 0} \langle f, \varphi_k \rangle \langle \varphi, \varphi_k \rangle = \sum_{k \geq 0} f_k \langle \varphi, \varphi_k \rangle$, in view of the preceding results we expect

$$(5.1) \quad \hat{f}_\varphi = \sum_{k \geq 0} \hat{f}_k \langle \varphi, \varphi_k \rangle,$$

to be an asymptotically efficient estimator. Before proceeding we need to introduce the extra condition that

$$(5.2) \quad \sum_{k \geq 0} \frac{1}{\rho_k} |\langle \varphi, \varphi_k \rangle| < \infty.$$

This condition ensures that φ is in the range of $(K^{-1})^*$. Let us write

$$(5.3) \quad \begin{aligned} \gamma &= (K^{-1})^* \varphi = V R^{-1} \left(\sum_{k \geq 0} \langle \varphi, \varphi_k \rangle \varphi_k \right) \\ &= \sum_{k \geq 0} \frac{1}{\rho_k} \langle \varphi, \varphi_k \rangle \varphi_{V,k}. \end{aligned}$$

Thanks to (2.10) and (5.2) the convergence in (5.3) is even pointwise.

To verify the asymptotic efficiency let us first observe that the optimal variance in the normal component of regular sequences of estimators equals

$$(5.4) \quad \tilde{\sigma}_\varphi^2(f) = \frac{\|\gamma\|^2}{I(\psi)}.$$

This can be shown by virtually the same method as employed in Section 3 for φ_0 .

Writing the decomposition in (4.2) for general φ_k (rather than for φ_0) as

$$(5.5) \quad \sqrt{n}(\hat{f}_k - f_k) = A_{n,k} + Q_{n,k} + R_{n,k},$$

we have

$$(5.6) \quad \begin{aligned} \sqrt{n}(\hat{f}_\varphi - f_\varphi) &= \sum_{k \geq 0} \langle \varphi, \varphi_k \rangle A_{n,k} + \sum_{k \geq 0} \langle \varphi, \varphi_k \rangle (Q_{n,k} + R_{n,k}) \\ &= S_{n,1} + S_{n,2}. \end{aligned}$$

As follows from (5.3),

$$(5.7) \quad \begin{aligned} S_{n,1} &= \frac{1}{\sqrt{n}} \sum_{i=1}^n \frac{1}{I(\psi)} \Lambda(\varepsilon_i) \sum_{k \geq 0} \frac{1}{\rho_k} \langle \varphi, \varphi_k \rangle \varphi_{V,k}(X_i) \\ &= \frac{1}{\sqrt{n}} \sum_{i=1}^n \frac{1}{I(\psi)} \Lambda(\varepsilon_i) \gamma(X_i), \end{aligned}$$

which means that $S_{n,1}$ is well defined. Exploiting the calculations in the proof of the Theorem in Section 4 it follows that $S_{n,2}$ is also well defined and that

$$(5.8) \quad \sum_{k \geq 0} \langle \varphi, \varphi_k \rangle (Q_{n,k} + R_{n,k}) = o_p(1).$$

Combination yields the following results.

Theorem. *Suppose that condition (5.2) is fulfilled in addition to the assumptions in Section 2. Then $\hat{f}_\varphi$ is an asymptotically efficient estimator of f_φ , i.e.*

$$(5.9) \quad \sqrt{n}(\hat{f}_\varphi - f_\varphi) \rightarrow_d \text{Normal} \left(0, \frac{\|\gamma\|^2}{I(\psi)} \right), \text{ as } n \rightarrow \infty,$$

6. Some comments on the conditions and improved regression estimation

6.1. The conditions. In the example of Section 1 the operator K itself is Hermitian and hence $V = I$ so that

$$(6.1) \quad \varphi_k(x) = \varphi_{V,k}(x) = \sqrt{2} \sin(k+1)\pi x, \quad 0 \leq x \leq 1, \quad k = 0, 1, \dots$$

It is immediate that conditions (2.5), (2.10), and (2.11) are fulfilled. It should be noted, however, that the function $\varphi \equiv 1$ on $[0, 1]$ doesn't satisfy (5.2). If the total weight is to be estimated, one should therefore employ a sufficiently smooth (near 0 and 1) approximation of this function.

Next suppose that the input function f satisfies $|\langle f, \varphi_k \rangle| = |f_k| \asymp k^{-s}$ for some $s > \frac{1}{2}$, and that $m(n) \asymp n^r$ for some $0 < r < \frac{1}{2}$. Then we have

$$(6.2) \quad \sqrt{n} \sum_{k > m} f_k^2 = O(n^{1/2+r(1-2s)}),$$

$$(6.3) \quad \sum_{k > m} |f_k| = O(n^{r(1-s)}), \quad s > 1,$$

and apparently conditions (2.13) and (2.14) are satisfied if

$$(6.4) \quad 0 < r < \frac{1}{2}, \quad s > \frac{1+2r}{4r}.$$

The conditions on the error density also appear to be usually fulfilled. A non-normal density that satisfies conditions (2.15) - (2.18) is, for instance, the logistic density

$$(6.5) \quad \psi(x) = \frac{e^{-x}}{(1 + e^{-x})^2}, \quad x \in \mathbb{R}.$$

In particular Λ' and Λ'' turn out to be bounded indeed.

6.2. Improving regression estimation. It should be noted that for the present results it is irrelevant whether the input estimator $\hat{f}_{(m)}$ attains the optimal MISE rate. It is not hard to see, however, that conditions on m and f ensuring this rate to be optimal are in general compatible with those in (2.12) - (2.14). In this discussion, however, we will allow the truncation index $M = M(n)$ of the traditional input estimator $\hat{f}_{(M)}$, that attains the optimal MISE rate, to tend to infinity at a different rate than the $m = m(n)$ used above.

The results in this paper regard the variances of the limiting normal distributions of the estimators, and not the variances or MSE's of these estimators themselves. It is clear, however, that

$$(6.6) \quad \text{MSE}(\hat{f}_0) = \text{Var} \hat{f}_0 = \frac{1}{n} \sigma_0^2(f),$$

and we conjecture that

$$(6.7) \quad \text{MSE}(\hat{\hat{f}}_0) = \mathbf{E}(\hat{\hat{f}}_0 - f_0)^2 = \frac{1}{n} \frac{1}{I(\psi)} + o\left(\frac{1}{n}\right).$$

This would imply that $\hat{\hat{f}}_0$ improves $\hat{f}_0$ also with respect to the MSE (at the level of constants). We similarly expect each of the $\hat{\hat{f}}_k$ to improve $\hat{f}_k$ regarding MSE, and eventually

$$(6.8) \quad \hat{\hat{f}}_{(M)}(y) = \sum_{k \leq M} \hat{\hat{f}}_k \varphi_k(y), \quad y \in \mathbf{Y},$$

to improve $\hat{f}_{(M)}$ with respect to MISE (at the level of constants). The actual calculations leading to (6.7) will differ from those in Section 4 and might be lengthy. Moreover, if theoretically $\hat{\hat{f}}_{(M)}$ would improve $\hat{f}_{(M)}$, it would be interesting to perform simulations for several nonnormal error distributions, to get an insight into the difference of the performance for finite sample sizes. All this is beyond the scope of this paper.

REFERENCES

- [1] Bickel, P.J., Klaassen, C.A.J., Ritov, Y. & Wellner, J.A. (1993). *Efficient and Adaptive Estimation for Semiparametric Models*. Johns Hopkins Univ. Press, Baltimore.
- [2] Debnath, L. & Mikusiński, P. (1999). *Hilbert Spaces with Applications*, 2nd ed. Academic Press, San Diego.
- [3] Goldenshluger, A. & Pereverzev, S.V. (2000). Adaptive estimation of linear functionals in Hilbert scales from indirect white noise observations. *Probab. Th. Rel. Fields* **118**, 169-186.
- [4] Hájek, J. & Šidák, Z. (1967). *Theory of Rank Tests*. Academia, Prague.
- [5] Ibragimov, I.A. & Hasminskii, R.Z. (1984). On nonparametric estimation of the value of a linear functional in Gaussian white noise. *Th. Probab. Appl.* **29**, 18-32.
- [6] Johnstone, I.M. & Silverman, B.W. (1990). Speed of estimation in positron emission tomography and related inverse problems. *Ann. Statist.* **18**, 251-280.
- [7] Kac, M. (1966). Can you hear the shape of a drum? *Amer. Math. Monthly* **73**, 1-23.

- [8] Kirsch, A. (1996). *An Introduction to the Mathematical Theory of Inverse Problems*. Springer, New York.
- [9] Kress, R. (1989). *Linear Integral Equations*. Springer, New York.
- [10] Klaassen, C.A.J., Lee, E.-J. & Ruymgaart, F.H. (2001). On efficiency of indirect estimation of nonparametric regression functions. *Contemporary Mathematics* **287**, 173-184.
- [11] Pfanzagl, J. (1994). *Parametric Statistical Theory*. De Gruyter, Berlin.
- [12] Riesz, F. & Sz.-Nagy, B. (1990). *Functional Analysis*. Dover, New York.
- [13] van Rooij, A.C.M., Ruymgaart, F.H. & van Zwet, W.R. (1999). Asymptotic efficiency of inverse estimators. *Th. Probab. Appl.* **44**, 826-844.
- [14] O'Sullivan, F. (1986). A statistical perspective on ill-posed inverse problems. *Statist. Sci.* **4**, 502-527.
- [15] van der Vaart, A.W. (1998). *Asymptotic Statistics*. Cambridge Univ. Press.

KORTEWEG-DE VRIES INSTITUTE FOR MATHEMATICS, UNIVERSITY OF AMSTERDAM, AMSTERDAM, THE NETHERLANDS

E-mail address: `chrisk@science.uva.nl`

DEPARTMENT OF MATHEMATICS AND STATISTICS, TEXAS TECH UNIVERSITY, LUBBOCK, TX 79409, USA

E-mail address: `elee@math.ttu.edu`

DEPARTMENT OF MATHEMATICS AND STATISTICS, TEXAS TECH UNIVERSITY, LUBBOCK, TX 79409, USA

E-mail address: `ruymg@math.ttu.edu`